

University of Groningen

## Foraging in a tidally structured environment by red knots (*Calidris canutus*)

van Gils, Jan A; Spaans, B; Dekinga, A; Piersma, T

*Published in:*  
Ecology

*DOI:*  
[10.1890/0012-9658\(2006\)87\[1189:FIATSE\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1189:FIATSE]2.0.CO;2)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2006

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

van Gils, J. A., Spaans, B., Dekinga, A., & Piersma, T. (2006). Foraging in a tidally structured environment by red knots (*Calidris canutus*): Ideal, but not free. *Ecology*, 87(5), 1189-1202. [https://doi.org/10.1890/0012-9658\(2006\)87\[1189:FIATSE\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1189:FIATSE]2.0.CO;2)

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.


### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

---

## HIGHLY SOCIAL FORAGER



Red Knots (*Calidris canutus*) mainly feed on hard-shelled bivalves, which they ingest whole. As bivalves live buried in the sediment, knots need to actually feed in order to “get a feel” for prey density. Nevertheless, as can be derived from the birds’ foraging behavior (patch selection, foraging intensity, and energy stores), the study shows that knots are very well informed about the spatial distribution of their food. As a highly social forager, these birds may have access to many forms of public information. However, knots do not always occupy the best patches, since, like all animals, they must bear the cost of traveling. We therefore conclude that Red Knots can be characterized as “ideal, nonfree foragers.” (Photo copyright Jan van de Kam, used by permission, all rights reserved.)

For more details see the paper “Foraging in a tidally structured environment by Red Knots (*Calidris canutus*): ideal, but not free,” by J.A. van Gils, B. Spaans, A. Dekinga, and T. Piersma, to be published in *Ecology* **87**(5), May 2006.